



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Designing and Synthesis of Site-Specific DNA Cleavage Reagents

Hai Liu^a, Xiqing Li^b, Bo Tan^a, Yang Zhao^a, Changde Lu^b & Yu-Fen Zhao^a

^a Tsinghua University, China

^b Chinese Academic of Sciences, China

Published online: 27 Oct 2010.

To cite this article: Hai Liu, Xiqing Li, Bo Tan, Yang Zhao, Changde Lu & Yu-Fen Zhao (2002) Designing and Synthesis of Site-Specific DNA Cleavage Reagents, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 177:8-9, 2085-2085, DOI: [10.1080/10426500213418](https://doi.org/10.1080/10426500213418)

To link to this article: <http://dx.doi.org/10.1080/10426500213418>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>



DESIGNING AND SYNTHESIS OF SITE-SPECIFIC DNA CLEAVAGE REAGENTS

Hai Liu,^a Xiqing Li,^b Bo Tan,^a Yang Zhao,^a Changde Lu,^b
and Yu-Fen Zhao^a
Tsinghua University, China^a and Chinese Academic
of Sciences, China^b

(Received July 29, 2001; accepted December 25, 2001)

In the past decade, studies on nucleic acids site-specific cleavage reagents, which consist of cleavage and recognition systems, had been received significant attention in the field of chemistry, biochemistry, and molecular biology. Previously it has been found that the dipeptide seryl-histidine (Ser-His) and related oligopeptides can cleave DNA, protein, and the ester p-nitrophenyl acetate (p-NPA) over wide ranges of pH and temperature.¹ Denaturing polyacrylamide gel electrophoresis (PAGE) of 50-end labeled DNA samples incubated with Ser-His reveals a pattern of two bands per nucleotide position, consistent with the generation of both 30-hydroxyl and 30-phosphate DNA cleavage fragments, as would be expected of phosphodiester hydrolysis by Ser-His. In order to confirm its hydrolysis mechanism and develop novel site-specific cleavage reagents, three different kinds of cleavage reagent consisted with recognition chain and cleavage active-site have been designed and synthesized. All of the recognition chains are made up of antisense oligonucleotide. The synthesis involved 10 steps. Further investigations on the cleavage mechanism and biological activities are current in progress.

REFERENCE

- [1] Li Yunsheng, Yu-Fen Zhao, Scott Hatfield, and Rong Wan, *Bioorganic & Medicinal Chemistry*, **8**, 2675–2680 (2000).

Address correspondence to Yu-Fen Zhao, Bioorganic Phosphorus Chemistry Laboratory, Department of Chemistry, School of Life Science and Engineering, Tsinghua University, Beijing 100084, P. R. China. E-mail: yfzhao@mail.tsinghua.edu.cn